

# Urine Mutagenicity as an Indicator of Exposure to Dietary Mutagens Formed During Cooking of Foods

by Robert S. U. Baker,\* Ian Darnton-Hill,\* Antonio M. Bonin,\* Andrew Arlauskas,\* Candida Braithwaite,\* Michael Wootton,<sup>†</sup> and A. Stewart Truswell<sup>‡</sup>

Studies were undertaken with individuals fed fried bacon meals to determine whether fruit or vegetables, ingested along with bacon, modified uptake and subsequent excretion of bacon mutagen(s). Urinary mutagenic activity was significant in those who had consumed bacon or mixed bacon/vegetable or bacon/fruit meals within the previous 2 to 3 hr period. Although urine activity varied by a factor of 4 among 15 subjects who consumed different meals, there was no evidence from this investigation that fruit or vegetables contributed to the inherent variability in total urinary mutagenic activity. However, some differences in excretion kinetics may be attributable to vegetable or fruit supplements in mixed meals.

## Introduction

Urinary excretion is one of the mechanisms the body has available to it for eradicating xenobiotics. Whether or not foreign molecules harm the body depends on their pharmacological activity, which in turn depends on tissue distribution, metabolic pathways in specific tissues, rates of metabolism, and on the balance between intake and elimination from the body.

Notwithstanding the problem of defining precise kinetics of absorption, distribution, etc., urinary excretion has been widely used in toxicology as an exposure barometer for potentially hazardous chemicals encountered in the workplace and for medicinal and other drugs (1,2).

As well as toxins, mutagenic substances may be detected in urine. Durston and Ames (3) demonstrated the mutagenicity of a urinary metabolite excreted as a glucuronide conjugate in urine of rats administered the carcinogen, 2-acetylaminofluorene. Benzidine, *o*-toluidine, *o*-tolidine, and aniline or their mutagenic metabolites were shown to occur in the urine of rats fed with

these aromatic amines (4). Mice administered the drug, metronidazole, excreted a mutagenic derivative, also detected in human urine (5). Furthermore, occupational exposure of humans to chemotherapeutic agents (6), anesthetic gases (7), epichlorhydrin (8), and some industrial chemicals (9,10) has apparently contributed to excess mutagenic activity detectable in urine. Thus, measurement of urinary activity has been proposed as a means of monitoring absorption of mutagens (11,12).

The search for mutagenic excretion products, which might be related to human cancer, has inevitably led to studies of "lifestyle factors" such as cigarette smoking (13), hairdye use (14), and coffee consumption (15). We previously reported detecting urine mutagenicity following fried pork or bacon meals (16). Since then Sousa, Nath, and Ong (17) have found a similar effect following beef meals. Both studies emphasize the absorption, distribution, and excretion of biologically detectable mutagenic substances. This finding is significant in view of the fact that nutritional components are considered to play an important role in induction of a number of human cancers (18).

In the present paper we examine the effect of other dietary constituents on urine mutagen excretion. Since vitamins, minerals, peroxidases, chlorophyll, etc. can inactivate or inhibit activation of pyrolysis mutagens (19), it is possible that similar components of vegetables and fruit, eaten along with fried meat, may directly

\*Department of Public Health Biology, School of Public Health and Tropical Medicine, University of Sydney, NSW, 2006, Australia.

<sup>†</sup>School of Food Technology, University of NSW, Kensington, NSW, 2033, Australia.

<sup>‡</sup>Human Nutrition Unit, Department of Biochemistry, University of Sydney, NSW, 2006, Australia.

also explain why there is no correlation between the amount of bacon eaten and the total urinary mutagenic activity/person/8 hr. A similar lack of correlation has been reported between the number of cigarettes smoked and mutagen levels in smokers' urines (25).

From this study, urine activity can be used as an indicator of exposure, but not as a dosimeter of the extent of exposure, to dietary mutagens. Thus, urinary excretion may be used to investigate inhibitory or protective factors in the diet, provided individual variation is taken into account. There is no evidence here that overall urinary clearance of bacon mutagen(s) was modified by fruit or vegetables, eaten with bacon, despite the evidence of inactivation and adsorption of pyrolysis mutagens by vegetable constituents (20,26).

This and similar studies point to the need to take into account individual variability, the contribution of dietary mutagens and of inhibitors of mutagenesis as potential confounding factors in using urine assays for monitoring environmental or occupational exposures.

## REFERENCES

1. Zielhuis, R. L. Biological monitoring. *Scand. J. Work Environ. Health* 4: 1-18 (1978).
2. Van Doorn, R., Leijdekkers, Ch. M., Bos, R. P., Brouns, R. M. E., and Henderson, P. Th. Detection of human exposure to electrophilic compounds by assay of thioether detoxication products in urine. *Ann. Occup. Hyg.* 24: 77-92 (1981).
3. Durston, W. E., and Ames, B. N. A simple method for the detection of mutagens in urine: studies with the carcinogen 2-acetylaminofluorene. *Proc. Natl. Acad. Sci. (U.S.)* 71: 737-741 (1974).
4. Tanaka, K. -I., Marui, S., and Mii, T. Mutagenicity of extracts of urine from rats treated with aromatic amines. *Mutat. Res.* 79: 173-176 (1980).
5. Connor, T. H., Stoeckel, M., Evrard, J., and Legator, M. S. The contribution of metronidazole and two metabolites to the mutagenic activity detected in urine of treated humans and mice. *Cancer Res.* 37: 629-633 (1977).
6. Falck, K., Gröhn, P., Sorsa, M., Vainio, H., Heinonen, E., and Holsti, L. R. Mutagenicity in urine of nurses handling cytostatic drugs. *Lancet* i: 1250-1251 (1979).
7. McCoy, E. C., Hankel, R., Rosenkranz, H. S., Giuffrida, J. G., and Bizzari, D. V. Detection of mutagenic activity in the urine of anesthesiologists: a preliminary report. *Environ. Health Perspect.* 21: 221-223 (1977).
8. Kilian, D. J., Pullin, T. G., Connor, T. H., Legator, M. S., and Edwards, H. N. Mutagenicity of epichlorohydrin in the bacterial assay system: evaluation by direct in vitro activity of urine from exposed humans and mice. *Mutat. Res.* 53: 72 (1978).
9. Falck, K., Sorsa, M., Vainio, H., and Kilpikari, I. Mutagenicity in urine of workers in rubber industry. *Mutat. Res.* 79: 45-52 (1980).
10. Schimberg, R. W., Skyttä, E., and Falck, K. Belastung von Eisingliesseriearbeitern durch mutagene polycyclische aromatische Kohlenwasserstoffe. *Staub-Reinh. Luft* 41: 421-424 (1981).
11. Dean, B. J., Ed. Report of the UKEMS Sub-committee on Guidelines for Mutagenicity Testing. Part II: Supplementary Tests. UKEMS, 1984, p203.
12. Vainio, H., Sorsa, M., and Hemminki, K. Biological monitoring in surveillance of exposure to genotoxicants. *Am. J. Ind. Med.* 4: 87-103 (1983).
13. Yamasaki, E., and Ames, B. N. Concentration of mutagens from urine by adsorption with the nonpolar resin XAD-2: cigarette smokers have mutagenic urine. *Proc. Natl. Acad. Sci. (U.S.)* 74: 3555-3559 (1977).
14. Burnett, C. M., Fuchs, C. M., and Corbett, J. F. Mutagenicity studies on urine concentrates from female users of dark hair color products. *Drug Chem. Toxicol.* 2: 283-293 (1979).
15. Aeschbacher, H. U., and Chapuis, C. Nonmutagenicity of urine from coffee drinkers compared with that from cigarette smokers. *Mutat. Res.* 89: 161-177 (1981).
16. Baker, R., Arlauskas, A., Bonin, A., and Angus, D. Detection of mutagenic activity in human urine following fried pork or bacon meals. *Cancer Letters* 16: 81-89 (1982).
17. Sousa, J., Nath, J., and Ong, T. Detection of urinary mutagenic activity following a fried beef meal. *Environ. Mutagen.* 6: 476 (1984).
18. Williams, G. M., Weisburger, J. H., and Wynder, E. L. Lifestyle and cancer etiology. In: *Carcinogens and Mutagens in the Environment*. Vol. I, Food Products (H. F. Stich, Ed.), CRC Press, Boca Raton, FL, 1982, pp. 53-64.
19. Kada, T. Desmutagens: an overview. In: *Carcinogens and Mutagens in the Environment*. Vol. II, Naturally Occurring Compounds (H. F. Stich, Ed.), CRC Press, Boca Raton, FL, 1982, pp. 63-67.
20. Henry, R. J., Cannon, D. C., and Winkelman, J. W. *Clinical Chemistry: Principles and Technics*, 2nd Ed. Harper & Row, Hagerstown, 1974, p. 552.
21. Gibson, J. F., Baxter, P. J., Hedworth-Whitty, R. B., and Gompertz, D. Urinary mutagenicity assays: a problem arising from the presence of histidine associated growth factors in XAD-2 prepared urine concentrates, with particular relevance to assays carried out using the bacterial fluctuation test. *Carcinogenesis* 4: 1471-1476 (1983).
22. Arimoto, S., Ohara, Y., Namba, T., Negishi, T., and Hayatsu, H. Inhibition of the mutagenicity of amino acid pyrolysis products by hemin and other biological pyrrole pigments. *Biochem. Biophys. Res. Commun.* 92: 662-668 (1980).
23. Hayatsu, H., Inoue, K., Ohta, H., Namba, T., Togawa, K., Hayatsu, T., Makita, M., and Wataya, Y. Inhibition of the mutagenicity of cooked-beef basic fraction by its acidic fraction. *Mutat. Res.* 91: 437-442 (1981).
24. Dolara, P., Caderni, G., Salvadori, M., Tringale, L., and Lodovici, M. Urinary mutagens in humans after fried pork and bacon meals. *Cancer Letters* 22: 275-280 (1984).
25. Jaffe, R. L., Nicholson, W. J., and Garro, A. J. Urinary mutagen levels in smokers. *Cancer Letters* 20: 37-42 (1983).
26. Kada, T., Kato, M., Aikawa, K., and Kiriya, S. Adsorption of pyrolysate mutagens by vegetable fibres. *Mutat. Res.* 141: 149-152 (1984).